

Prevention and Management of Diabetic Foot Disease



Dr Samer Alsabbagh

Rebecca Gardner

Dr Chantal Kong

Caroline Leith

Dr Pawan Pusalkar

Carolyn Wareham

Format


- Background & Epidemiology
- Case studies
- Standards of Care
- Guidelines and Referral Care Pathways
- Management of the diabetic foot
- Summary
- Question time

Epidemiology of the Diabetic Foot

- Leading cause of all non traumatic lower limb amputation (40-60%)
- Commonest cause of hospital bed occupancy (most common cause of hospital admission amongst diabetes patients, NaDIA)
- 85% are preceded by foot ulceration
- Lower limb amputations \uparrow x 15 in diabetes
- > 50% require amputation of other limb within 3-5 years
- 50% patients die within 5 years



“Are diabetes-related wounds and amputations worse than cancer?”¹



1. Adapted from Armstrong DG, Wrobel J, Robbins JM. Int Wound J. 2007 Dec; 4(4): 286-7.

Cost of poor diabetic foot care to NHS


In 2010-11, the NHS in England spent an estimated **£639 million-£662 million, 0.6–0.7% of its budget**, on diabetic foot ulceration and amputation

	Lower estimate	Upper estimate
Primary, community and outpatient care	£306,508,970	£323,062,601
Accident and emergency		£849,278
Inpatient care – ulceration	£213,151,916	£213,151,916
Inpatient care – amputation	£43,546,901	£48,896,732
Post-amputation care	£75,807,423	£75,807,423
Total	£639,015,210	£661,767,953

Estimated cost of ulceration and amputation in people with diabetes, England, 2010–11

**£1.75-
£1.81
million
a day!**

Hospitalisation represents the greatest proportion of overall costs¹




Variation in amputation rates 'shocking' (BBC News 2010, 2012)

- Department of Health data reveals the rate of major amputations in the South West, at 3 in 1000, almost twice the rate in the South East
- Are we doing any better in Hertfordshire?

<http://www.bbc.co.uk/news/health-17270379>[17/08/2012 16:49:30]

<http://www.bbc.co.uk/news/health-19050684>[17/08/2012 16:48:14]



Diabetes Footcare Activity Profile August 2016

(www.gov.uk/phe, www.ncvin.org.uk)

Key facts	Herts Valleys CCG	East & North Herts CCG	England
Total episodes of inpatient care for diabetic foot disease	1,776	1,241	167,224
Annual episodes of care for diabetic foot disease per 1,000 adults with diabetes	24.6	16.6	19.8
Total nights in hospital due to diabetic foot disease	13,006	11,731	1,357,714
Annual nights in hospital for diabetic foot disease per 1,000 adults with diabetes	180.1	156.6	161.0
Episodes of care where an amputation is performed on those with diabetes	212	119	22,109
Annual amputations per 1,000 adults with diabetes	2.9	1.6	2.6

Diabetes Footcare Activity Profile August 2016

(www.gov.uk/phe, www.ncvin.org.uk)

Key facts	Herts Valleys CCG	East & North Herts CCG	England
Total episodes of inpatient care for diabetic foot disease	1,776	1,241	167,224
Annual episodes of care for diabetic foot disease per 1,000 adults with diabetes	24.6	16.6	19.8
Total nights in hospital due to diabetic foot disease	13,006	11,731	1,357,714
Annual nights in hospital for diabetic foot disease per 1,000 adults with diabetes	180.1	156.6	161.0
Episodes of care where an amputation is performed on those with diabetes	212	119	22,109
Annual amputations per 1,000 adults with diabetes	2.9	1.6	2.6

Why the variation in foot outcomes?

Root cause analysis

- What structures and systems are in place locally? (commissioning/organisation)
- Are there clear guidelines/protocols for
 - referral routes/care pathways?
 - process of care?
- Do patients & health care professionals know them, how and where to access them?
- What are the current resources (podiatry, primary and secondary care, inpatients)
- Are all patients being risk assessed appropriately?
- Are all patients at risk being followed up as required?
- Are the appropriate review/discharge care planning arrangements in place? (Continuity of care)
- Optimal communication between HCPs involved (Use of different IT system)?


Case studies

What level of care should the diabetic patient with a foot problem expect?

What can **you** do to improve your diabetic patients' footcare?

Quality standards

- Putting Feet First 2003
- NICE 2004 CG10 (Type 2 Diabetes foot problems: Prevention and Management of Foot problems)



Quality standards

- Putting Feet First 2003
- NICE 2004 CG10 (Type 2 Diabetes foot problems: Prevention and Management of Foot problems)
- NICE CG15 2004 (Diagnosis and management of Type 1 diabetes in children, young people and adults)

NHS
National Institute for
Clinical Excellence

Prevention and
management of foot
problems in people with

National Institute for
Clinical Excellence

Type 1 diabetes in children and young people

Understanding NICE guidance – information for
the families and carers of children with type 1
diabetes, young people with type 1 diabetes,
and the public

2010
The recommendation to re-test for coeliac disease
every 3 years has been removed.

Some other changes have been made in line with
changes in other guidance.

In this document changes are marked with black strikethrough.
Details of all changes can be found at
www.nice.org.uk/guidance/CG15/NICEGuidance/ChangesApr2010

July 2004


Corporate member of
Plain English Campaign.
Committed to clearer communication.

197




Quality standards

- Putting Feet First 2003
- NICE 2004 CG10 (Type 2 Diabetes foot problems: Prevention and Management of Foot problems)
- NICE CG15 2004 (Diagnosis and management of Type 1 diabetes in children, young people and adults)
- **NICE CG119 March 2011 (Diabetic foot problems: Inpatient management of diabetic foot problems)**



Quality standards

- Putting Feet First 2003
- NICE 2004 CG10 (Type 2 Diabetes foot problems: Prevention and Management of Foot problems)
- NICE CG15 2004 (Diagnosis and management of Type 1 diabetes in children, young people and adults)
- NICE CG119 March 2011 (Diabetic foot problems: Inpatient management of diabetic foot problems)
- **NICE CG19 Aug 2015, updated Jan 2016 (Diabetic foot problems: prevention and management)**




Commissioners and service providers should ensure that the following are in place:

- A **foot protection service** for preventing diabetic foot problems, and for treating and managing diabetic foot problems in the community
- A **multidisciplinary foot care service** for managing diabetic foot problems in hospital and in the community that cannot be managed by the foot protection service
- **Robust protocols and clear local pathways** for the **continued and integrated care** of people across all settings, including **emergency care and general practice**. The protocols should set out the relationship between the foot protection service and the multidisciplinary foot care service
- Regular reviews of treatment and patient outcomes, in line with the [National Diabetes Foot Care Audit \(NDFA\)](#)

Reducing the risk of developing a diabetic foot problem

- Education, Education, Education.....
(patients, carers, HCPs)
- On-going care: annual review & recall
- Detection of risk factors for ulceration
- Classification of foot risk
- Refer early to Community Podiatry & MDT foot clinics



How frequently should you assess your diabetic patients' feet?

- At time of diagnosis and at least annually thereafter
- If any foot problems arise
- On any admission to hospital, and if there is any change in their status while they are in hospital

Assessing the risk of developing a diabetic foot problem

Remove the patient's shoes, socks, bandages and dressings, and examine both feet for evidence of the following risk factors:

- Neuropathy (use a 10 g monofilament as part of a foot sensory examination)
- Limb ischaemia
- Ulceration
- Callus
- Infection and/or inflammation
- Deformity
- Gangrene
- Swelling (Charcot arthropathy)



Factors associated with foot ulcer

- Previous ulcer/amputation
- Neuropathy
 - Sensorimotor
- Trauma
 - Poor footwear
 - Walking barefoot
 - Falls/accidents
 - Objects inside shoes
- Biomechanics
 - Limited joint mobility
 - Bony prominences
 - Foot deformity/osteoarthropathy
 - Callus
- Peripheral vascular disease
- Socio-economic status
 - Low social position
 - Poor access to healthcare
 - Non compliance/neglect
 - Poor education

Foot Risk stratification for the patient with diabetes (1)

- Low risk (normal sensation, palpable pulses, +/- callus)
- Moderate (increased) risk
- High risk
- Active & the Urgent/Emergency diabetic foot

Active diabetic foot problem

- ulceration **or**
 - spreading infection
-
- **Refer urgently to the MDT Specialist Foot clinic**

This is an emergency

- critical limb ischaemia **or**
 - gangrene **or**
 - suspicion of an acute Charcot arthropathy, or an unexplained hot, red, swollen foot with or without pain
-
- **Refer urgently to AAU**

PUTTING FEET FIRST

Annual Foot Review for everyone with diabetes over 12 years old

How to do an annual foot check:

- Remove shoes and socks/ stockings
- Test foot sensations using 10g monofilament or vibration with a tuning fork
- Palpate foot pulses
- Inspect for any deformity
- Inspect for significant callus
- Check for signs of ulceration
- Ask about any previous ulceration
- Inspect footwear
- Ask about any pain
- Tell patient how to look after their feet and provide written information
- Tell patient their risk status and what it means. Explain what to look out for and provide emergency contact numbers.

ADVISE THE PATIENT TO:

- Check their feet every day
- Be aware of loss of sensation
- Look for changes in the shape of their foot
- Not use corn removing plasters or blades
- Know how to look after their toenails
- Wear shoes that fit properly
- Maintain good blood glucose control
- Attend their annual foot review

IDENTIFICATION OF FOOT RISK STATUS AND THE ACTION TO TAKE

LEVEL OF RISK

ACTION

ACTIVE

- Ulceration or spreading infection or critical limb ischaemia (severe peripheral arterial disease) or gangrene or suspicion of acute Charcot foot or an unexplained hot, red, swollen foot with or without pain.

- Rapid referral (within one working day) to the Foot Protection Service (FPS) or the multidisciplinary foot team, for triage within one further working day.
- Assess feet and lower limbs, then agree a tailored treatment plan.
- Provide written and verbal education with emergency contact numbers.
- Refer for special intervention if/ when required.
- Liaise with other healthcare professionals eg GP as necessary.

HIGH

- Previous ulceration or previous amputation or on renal replacement therapy (dialysis or transplant) or neuropathy (loss of sensation) and lower limb peripheral arterial disease together or neuropathy (loss of sensation) in combination with callus and/ or deformity* or lower limb peripheral arterial disease in combination with callus and/ or deformity*.

- Refer to a specialist podiatrist or member of the foot protection service (FPS) and request an assessment within 2-4 weeks.
- Thereafter they should be assessed every 1-2 months if there is no immediate concern. This is in addition to their annual assessment. Both assessments should be carried out by a specialist podiatrist or a member of the FPS.
- Assess feet and lower limbs, then agree a tailored treatment plan.
- Provide written and verbal education with emergency contact numbers.
- Refer for special intervention if/ when required.
- Liaise with other healthcare professionals eg GP as necessary.

MODERATE

- Deformity* or neuropathy (loss of sensation) or lower limb peripheral arterial disease.

- Refer to a specialist podiatrist or member of the foot protection service (FPS) and request an assessment within 6-8 weeks.
- Thereafter they should be assessed every 3-6 months in addition to their annual assessment, by a specialist podiatrist or a member of the FPS.
- Assess feet and lower limbs, then agree a tailored treatment plan.
- Provide written and verbal education with emergency contact numbers.
- Refer for special intervention if/ when required.
- Liaise with other healthcare professionals eg GP as necessary.

LOW

- No risk factors, as listed above, present.
- Callus alone is considered low risk.

- Annual screening by a suitably trained Healthcare Professional.
- Agree self management plan.
- Provide written and verbal education with emergency contact numbers.

Record risk status and inform patient of their risk status and what it means.

*A change in foot shape that results in difficulty in fitting a standard shoe, as assessed by the practitioner.


These risk categories relate to the use of the SCI-DC foot risk stratification tool and NICE guidance (NG19, 2015).

Produced by the Scottish Diabetes Foot Action Group



DIABETES UK
CARE. CONNECT. CAMPAIGN.

Managing a diabetic foot problem



Key priorities are to:

- Treat any infection that is present
- Treat vascular disease, if present
- Alleviate pressure to aid healing (off-loading)
- Achieve good metabolic (glycaemic) control and control of risk factors for cardiovascular disease (smoking, dyslipidaemia)

Emergency referral

- **Refer within 24 hrs to multidisciplinary specialist foot care team (MDT Foot clinics) if**
 - new ulcer
 - new swelling
 - new discolouration
 - no critical ischaemia (needs urgent vascular review)

Management of active foot ulcers (1)

- If clinical signs of active infection (redness, pain, swelling, discharge), give intensive systemic antibiotic therapy
 - Flucloxacillin 1g qds PO for 7-14 days
 - Doxycycline 100 mgs bd if penicillin allergic
- Use wound dressings that best match clinical experience, patient preference, site of the wound (no strong evidence). Consider cost of dressing
- Regular monitoring & dressing change of wounds
- Wound debridement (by specialist podiatrists or vascular surgeons)
- Plain Foot Xray if wound persistent and deep with oedema, suspicious of osteomyelitis
- Refer urgently to MDT Foot clinics **and** Community Podiatry
- Admit if moderate to severe infection especially if patient septic or evidence of critical ischaemia

Management of diabetic foot ulcers (2)

For a diabetic foot ulcer to heal, the following conditions must be satisfied


- Arterial inflow is adequate
- Infection is treated appropriately
- Pressure is removed from the wound and the immediate surrounding area

The most common cause of non-healing of neuropathic foot ulcers is **the failure to remove pressure** from the wound and immediate surrounding area



People with foot care emergencies and foot ulcers

- Foot care emergency (new ulcer, swelling, discolouration)
- Refer to multidisciplinary foot care team within 24 hrs
- Investigate & treat vascular insufficiency
- Initiate & supervise wound management
 - appropriate dressings & debridement
 - systemic antibiotics for cellulitis or/and bone infection
- Specialist footwear, orthotics & casts
- Optimise metabolic control, glucose levels, control cardiovascular risk factors



Multi-disciplinary team

Antibiotic pharmacist

Primary care team

Diabetes nurse

Splints/casting

Vascular surgeon

Microbiologist

Orthopaedic surgeon

Orthotist

Physician

Podiatrist



Diabetic Foot Osteomyelitis (DFO)

- **Diabetic foot Osteomyelitis** is a common problem, found in 20% overall to >60% (severe) diabetic foot infections – increase risk of LE amputation (up to 23)
- **Suspect if:**
 - long wound duration, recurrent infection
 - wound deep >3 mm, large > 2 cm, bony prominence visible
 - bone/joint - “sausage” toe

Probe to bone is useful if done and interpreted correctly

- **Blood tests:** WBC, CRP, PCT, ? Biomarkers

(No new serology tests yet found useful)

- **Imaging**

- Plain foot X-ray is the first test, limited sensitivity (early) & Specificity (late) - ?
- Repeat in 6-8 weeks if required
- if advanced imaging needed, MRI current best, marrow oedema
- Newer Imaging (SPECT/PET/CT/MRI) promising

Management of Diabetic Foot Osteomyelitis (DFO)

- Management is medical if detected and treated early
- Infection typically contiguous spread from soft tissue
- For wound culture, tissue specimens should be obtained by scraping the base of the ulcer with a scalpel or curette
- Oral antibiotic Rx probably as good as parenteral

Microbiology, pathophysiology:

- **S aureus predominant, Coag negative Staph,**
 - Beta-hemolytic streptococci (group A, B, and others)
 - **Gram negative Rod (polymicrobial),**
 - Anaerobic organisms
-
- Clindamycin; Daptomycin; Rifampicin especially effective for bone infection; others Fluoroquinolones;

People with Charcot osteoarthropathy

- Immediate referral to multidisciplinary foot care team
- Immobilise affected joint(s)
- Longstanding management is offloading
- No pharmacological cure/treatment
- BUT closely correlated with glycaemic control



Total Contact Cast

Diabetes Foot care Pathway in Hertfordshire – How do we compare with NICE?

- **Currently NICE incompliant for new foot ulcers and urgent foot problems** (all patients should be referred to MDT foot team/clinics)
- **BUT** major capacity issue for MDT foot clinics across all trusts
- Absence of a dedicated inpatient MDT specialist Footcare team (complete absence of inpatient podiatry at WHHT!!)
- Patients are being referred too late (NDFA 2016)
- ?Risk stratification of patients

Summary

- Significant proportion of amputations are preventable (but not all....)
- Early identification & treatment intervention is vital
- Management of the diabetic foot is complex and requires a multidisciplinary approach
- Refer early to the specialist podiatry team (Foot protection team) & MDT Diabetes Specialist Team
- Identify and treat any infection after a deep wound swab (NB Diabetic wounds may not always look actively infected)
- Regular foot checks vital
- Education of HCPs, patients & carers is essential
- Regular follow-up surveillance and education for all patients especially for moderate and high risk groups

Key messages/learning points

- Examine your patient's feet & risk stratify
- Recognise severity of the acute foot problem, especially infection & critical ischaemia
- Arrange appropriate investigations including plain Foot Xray
- Suspect underlying osteomyelitis if persistent non-healing ulcer
- Suspect acute Charcot's arthropathy in any unexplained leg/foot swelling
- Refer urgently to MDT Specialist Foot clinic if new ulcer
- Low threshold for antibiotics

Questions?

Discussion: How do we compare with NICE?

- Community support
- Foot education (prevention – patients & carers)
- Vulnerable patients
- Referral & access
- Process of care (assessment, antibiotics, communication with GPs & community)
- Multidisciplinary team
- Staffing levels v/s caseload
- Resource issues
- In-patients
- Education of HCPs